

## DESIGN AND SYNTHESIS OF TWO BICYCLO[3.3.1]NONANE-STEROID DERIVATIVES

Lauro Figueroa <sup>a\*</sup>, Alejandra Garcimarerero <sup>b</sup>, Rolando Garcia <sup>a</sup>, Francisco Diaz <sup>c</sup>,  
Marcela Rosas <sup>d</sup>, Virginia Mateu <sup>d</sup>, Maria Lopez <sup>a</sup>, Lenin Hau <sup>a</sup>, Tomas Lopez <sup>a</sup>,  
Abelardo Camacho <sup>e</sup>, Yaritza Borges <sup>a</sup>

<sup>a</sup>Faculty of Chemical-Biological Sciences, University Autonomous of Campeche,  
Agustín Melgar ave., Colonia Bellavista 24039, Campeche, Mexico

<sup>b</sup>Faculty of Medicine, University Veracruzana, Medicos and Odontologos str., Xalapa 91010, Veracruz, Mexico

<sup>c</sup>National School of Biological Sciences, National Polytechnic Institute,

Prolacion Carpio and Plan de Ayala str., Colonia Santo Tomas 11340, Mexico

<sup>d</sup>Faculty of Nutrition, University Veracruzana, Medicos and Odontologos str., Xalapa 91010, Veracruz, Mexico

<sup>e</sup>Faculty of Medicine and Nutrition, University Juarez of Durango State, Constitución str., Durango 34000, Mexico

\*e-mail: [lfiguero@uacam.mx](mailto:lfiguero@uacam.mx)

**Abstract.** Several studies for the synthesis of bicyclo[3.3.1]nonane analogues have been reported, however, there is little information on the preparation of bicyclo[3.3.1]nonane-steroid derivatives. In this way, the aim of this study was to synthesize two steroid-bicyclo[3.3.1]nonane analogues (**11** or **12**) from either estradiol or estrone using some reactions such as etherification, addition, nucleophilic substitution and cyclization. The chemical structure was evaluated through NMR spectroscopic analysis. The results showed higher yield for **11** compared with **12**. It is noteworthy, that the reagents used in this investigation are not expensive and do not require special conditions for handling.

**Keywords:** bicyclo, steroid, estrone derivative, synthesis, NMR.

Received: 27 December 2019/ Revised final: 10 March 2020/ Accepted: 14 March 2020

---