

ISOMERIZATION OF LIMONENE ON ZEOLITE-CONTAINING CATALYSTS BASED ON KAOLIN

Lyubov Patrylak^{ID a,b*}, Stepan Zubenko^{ID a}, Serhy Konovalov^{ID a}, Anzhela Yakovenko^{ID a},
Volodymyr Povazhnyi^{ID a}, Oleksandra Pertko^{ID a}, Yulia Voloshyna^{ID a},
Oleksandr Melnychuk^{ID a}, Mykhailo Filonenko^{ID a,c}

^a*V.P. Kukhar Institute of Bioorganic Chemistry and Petrochemistry of National Academy of Sciences of Ukraine,
1, Murmanska str., Kyiv 02094, Ukraine*

^b*National Technical University of Ukraine "Igor Sikorskyi Kyiv Polytechnic Institute",
37, Peremohy ave., Kyiv 03056, Ukraine*

^c*National Pedagogical Dragomanov University, 9, Pyrogova str., Kyiv 01601, Ukraine*
^{*}*e-mail: lkpg@ukr.net*

Abstract. The aim of the work was to study the isomerization of limonene on zeolite-containing biporous acid catalysts based on kaolin. Zeolite catalysts were synthesized from Ukrainian kaolin and characterized by using XRD, XRF, DTA/TG, IR-spectroscopy, low-temperature nitrogen adsorption, and pyridine sorption. Micro-mesoporous materials isomerize limonene at 160°C. The main reaction product on acid catalysts was terpinolene, while the original metakaolin microsphere catalyzes mainly the limonene condensation. The maximum yield of isomers is of 60–65% at 80–90% conversion. The obtained results show that the studied samples do not have a significant accumulation of carbonaceous deposits because limonene has high solubility, which helps to remove intermediate products of transformation from the surface of the samples.

Keywords: kaolin, zeolite, limonene isomerization, terpinolene yield.

Received: 20 June 2022/ Revised final: 08 September 2022/ Accepted: 14 September 2022
