

## MILD ACIDIC CHARCOAL CATALYZED SYNTHESIS OF 3,4-DIHYDROPYRIMIDIN-2(1H)-ONE/-THIONE DERIVATIVES

Rajendra Patil<sup>ID</sup><sup>a</sup>, Jagdish Chavan<sup>ID</sup><sup>a</sup>, Shivnath Patel<sup>ID</sup><sup>a</sup>, Vaishali Shinde<sup>b</sup>, Anil Beldar<sup>ID</sup><sup>a\*</sup>


<sup>a</sup>*Department of Chemistry, Poojya Sane Guruji Vidya Prasarak Mandal's Sajan Isan Patil Arts, Girdhar Barku Patel Science and Shahada Taluka Kharedi Vikri Sangh Commerce College, Khetia Road Shahada, Nandurbar 425409, India*

<sup>b</sup>*Department of Chemistry, Savitribai Phule Pune University, Ganeshkhind Road, Pune 411007, India*  
*\*e-mail: dragbeldar@gmail.com; phone: (+91) 94 20 134 233*

**Abstract.** A mild catalyst system with comparative reduction in amount of catalyst was demonstrated. The multicomponent synthesis of 3,4-dihydropyrimidin-2(1H)-ones and -thiones using acetic acid supported on activated charcoal as a mild acid catalyst in ethanol under both conventional as well as microwave irradiation conditions has been achieved. The obtained catalyst system is more efficient under microwave irradiation than under conventional conditions with shorter reaction times (3-9 min) and excellent yields (78-94 %).

**Keywords:** Biginelli reaction, acetic acid, acidic charcoal, microwave irradiation.

*Received: 20 September 2022/ Revised final: 18 November 2022/Accepted: 20 November 2022*



**Figure S1. IR Spectrum of 5-(Ethoxycarbonyl)-6-methyl-4-phenyl-3,4-dihydropyrimidin-2(1H)-one (4a).**




Figure S2. IR Spectrum of 5-(ethoxycarbonyl)-6-methyl-4-(4-nitrophenyl)-3,4-dihydropyrimidin-2(1H)-one (4c).




Figure S3. IR Spectrum of 5-(ethoxycarbonyl)-4-(4-methoxyphenyl)-6-methyl-3,4-dihydropyrimidin-2(1H)-one (4d).




Figure S4. IR Spectrum of 5-(ethoxycarbonyl)-6-methyl-4-(4-chlorophenyl)-3,4-dihydropyrimidin-2(1H)-thione (4h).






Figure S5. LCMS spectrum of 5-(ethoxycarbonyl)-6-methyl-4-phenyl-3,4-dihydropyrimidin-2(1H)-one (4a).



**Figure S6. LCMS spectrum of 5-ethoxycarbonyl-6-methyl-4-(4-hydroxyphenyl)-3,4-dihydropyrimidin-2(1H)-one (4b).**



**Figure S7. LCMS spectrum of 5-ethoxycarbonyl-4-(phenyl)-6-methyl-3,4-dihydropyrimidin-2(1H)-thione (4f).**




Figure S8. <sup>1</sup>H NMR spectrum of 5-(ethoxycarbonyl)-6-methyl-4-phenyl-3,4-dihydropyrimidin-2(1*H*)-one (4a).







Figure S9. <sup>1</sup>H NMR spectrum of 5-ethoxycarbonyl-6-methyl-4-(4-hydroxyphenyl)-3,4-dihydropyrimidin-2(1*H*)-one (4b).



**Figure S10.**  $^1\text{H}$  NMR spectrum of 5-(ethoxycarbonyl)-4-(n-propyl)-6-methyl-3,4-dihydropyrimidine-2(1*H*)-one (**4e**).



**Figure S11.**  $^1\text{H}$  NMR spectrum of 5-(Ethoxycarbonyl)-6-methyl-4-(4-chlorophenyl)-3,4-dihydropyrimidin-2(1*H*)-thione (4h).



**Figure S12.** <sup>1</sup>H NMR spectrum of 5-ethoxycarbonyl-4-(n-propyl)-6-methyl-3,4-dihydropyrimidin-2(1H)-thione (4j).