

CHEMICAL PERSPECTIVE ON RADIONUCLIDE EMISSIONS AS ATMOSPHERIC CONTAMINANTS FROM NUCLEAR REACTORS

Sergey Travin ^{a*}, Gheorghe Duca ^b, Oleg Gromov ^c

^a*N.N. Semenov Federal Research Centre for Chemical Physics of the Russian Academy of Sciences,
4, Kosygina str., Moscow 119991, Russian Federation*

^b*Institute of Chemistry of Moldova State University, 3, Academiei str., Chisinau, MD2028, Republic of Moldova*

^c*High Technology Research Institute of Inorganic Materials, A.A. Bochvar,
5a, Rogova str., Moscow 123098, Russian Federation*

**e-mail: travinso@yandex.ru*

Abstract. This study investigates the radionuclide emissions from nuclear power plants (NPPs), focusing on the gas-aerosol emissions from reactors such as VVER-440. The radionuclide composition of these emissions is analysed to determine the biological hazards they pose, particularly focusing on isotopes such as tritium and radiocarbon. The research highlights the patterns of radionuclide accumulation in fuel assemblies using "rank-size" coordinates, which provide a more visual and informative method compared to traditional atomic weight dependency analyses. The study introduces the Zipf-Mandelbrot type rank distributions and proposes that the emissions can be effectively monitored by measuring only a few key radionuclides, thereby simplifying the process. This method is validated across different reactor types, indicating its broad applicability. The research underscores the potential of using these patterns for more efficient regulation and control of radionuclide emissions from NPPs.

Keywords: radionuclide, NPP gas-aerosol emission, Zipf distribution, rank distribution, radioactive emission, fingerprint criterion.

Received: 25 September 2024/ Revised final: 19 December 2024/ Accepted: 20 December 2024
